

Junro Sato · Kiyoshi Baba

The chromatic number of the complementary graph to a simple graph associated with a commutative ring

Received: 1 November 2013 / Accepted: 9 February 2014 / Published online: 13 March 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract Let R be a Noetherian ring. We denote by $G(R)$ the simple graph whose vertices are elements of R and in which two distinct vertices x and y are joined by an edge if $x - y$ is a zero-divisor of R . Let $\overline{G}(R)$ be the complementary graph to $G(R)$ and $\overline{\chi}(R)$ be the chromatic number of the graph $\overline{G}(R)$. In this paper, we determine the chromatic number $\overline{\chi}(R)$. Let P_1, \dots, P_t be all maximal prime divisors of (0) . If $\overline{\chi}(R)$ is finite, then

$$\overline{\chi}(R) = \min\{|R/P_i|; i = 1, 2, \dots, t\}$$

where $|R/P_i|$ denotes the number of the set R/P_i .

Mathematics Subject Classification 05C15 · 13E05

المخلص

لتكن R حلقة نوثيرية. نرمز بـ $G(R)$ للبيان البسيط الذي تكون رؤوسه عناصر R ويتم فيه وصل رأسين مختلفين x و y بواسطة حافة إذا كان $x - y$ قاسماً للصفر في R . ليكن $\overline{G}(R)$ البيان المتكامل لـ $G(R)$ و $\overline{\chi}(R)$ العدد اللوني للبيان $\overline{G}(R)$. نحدد في هذه الورقة العدد اللوني $\overline{\chi}(R)$. لتكن P_1, \dots, P_t جميع القواسم الأولية الأعظمية لـ (0) . إذا كان $\overline{\chi}(R)$ منتهياً، فإن

$$\overline{\chi}(R) = \min\{|R/P_i|; i = 1, \dots, t\}$$

حيث يرمز $|R/P_i|$ لعدد عناصر المجموعة R/P_i .

First, we collect the basic notions and results of graph theory for later use. For a simple graph G , $V(G)$ denotes the set of vertices of G and $E(G)$ denotes the set of edges of G . We color the vertices of G so that no two joined vertices have the same color. If we color the vertices, we call it a coloring of G . The chromatic number $\chi(G)$ of the graph G is the minimum number of colors needed to color G .

Let C be a non-empty subset of $V(G)$. We call C a clique of G if every pair of two distinct elements of C is joined by an edge. The clique number $C(G)$ of G is the number of vertices in the largest clique of G .

For a set S , $|S|$ denotes the number of element of S .

Our notation is standard and for unexplained terms, our general reference to commutative algebra is [1] and [5], and our general reference to graph theory is [2].

J. Sato (✉)

Department of Mathematics, Faculty of Education, Kochi University, 2-5-1 Akebono-cho, Kochi 780-8520, Japan
E-mail: junro@kochi-u.ac.jp

K. Baba

Department of Mathematics, Faculty of Education and Welfare Science, Oita University, Oita 870-1192, Japan
E-mail: baba@cc.oita-u.ac.jp



Springer

We express our hearty gratitude to the referees for helpful comments on our paper.

Lemma 1 (cf. [6, Lemma 3]) *The inequality $\chi(G) \leq \chi(G)$ holds.*

The symbol \coprod denotes the disjoint union of sets.

Lemma 2 (cf. [6, Lemma 4]) *Let V_1, V_2, \dots, V_t be non-empty subsets of $V(G)$. Let*

$$V(G) = V_1 \coprod V_2 \coprod \dots \coprod V_t.$$

be a disjoint union of $V(G)$ such that no pair of two distinct elements of V_i is joined by an edge for $i = 1, 2, \dots, t$. Then $\chi(G) \leq t$.

Let G_1 and G_2 be two simple graphs. We say that G_1 is a subgraph of G_2 if the following conditions hold: (1) $V(G_1) \subset V(G_2)$, (2) $E(G_1) \subset E(G_2)$.

The proof of the following lemma is obvious, so we omit it.

Lemma 3 *If G_1 is a subgraph of G_2 , then $\chi(G_1) \leq \chi(G_2)$.*

Let G be a simple graph. We define the complementary graph \overline{G} of G to be a graph satisfying the following conditions:

- (1) $V(\overline{G}) = V(G)$.
- (2) Let x and y be two distinct vertices of \overline{G} . Then, x and y are joined by an edge in \overline{G} if and only if x and y are not joined by an edge in G .

Let R be a commutative ring with the identity element. An element x is called a zero-divisor of R if there exists a non-zero element y of R such that $xy = 0$. We denote by $Z(R)$ the set of zero-divisors of R . We consider the simple graph $G(R)$ whose vertices are elements of R and in which two distinct vertices x and y are joined by an edge if $x - y$ is in $Z(R)$. Let $\chi(R)$ denote the chromatic number of the graph $G(R)$ and let $V(R)$ denote the set of vertices of $G(R)$.

$\overline{G}(R)$ denotes the complementary graph to $G(R)$ and $\overline{\chi}(R)$ denotes the chromatic number of the graph $\overline{G}(R)$.

In our previous paper [6], we have shown that if $\chi(R)$ is finite, then R is an integral domain or a finite ring. On the other hand, even if $\overline{\chi}(R)$ is finite, R is not necessarily a finite ring as the following example shows.

Before we proceed to Example 4, we collect some notions and results of commutative algebra for later use.

Let N be an R -module. For an element x of N , $\text{Ann}_R(x) = \{c \in R; cx = 0\}$ is called the annihilator ideal of x . We say a prime ideal P of R is an associated prime ideal of N if there exists an element x of N such that $P = \text{Ann}_R(x)$. $\text{Ass}_R(N)$ denotes the set of all associated prime ideals of N . We consider the case $N = R$. If R is Noetherian, then $\text{Ass}_R(R)$ consists of prime divisors of (0) in the primary decomposition of (0) ([5, (8.E) Lemma and (8.G) Theorem 11]).

Example 4 Let \mathbf{Z} be the ring of integers and set $\mathbf{F}_2 = \mathbf{Z}/2\mathbf{Z}$. Let $\mathbf{F}_2[S, T]$ be a polynomial ring over \mathbf{F}_2 in indeterminates S and T . Let s and t be the residue classes of S and T in $\mathbf{F}_2[S, T]/(ST, S^2)$ respectively. Then, $\mathbf{F}_2[s, t] = \mathbf{F}_2[S, T]/(ST, S^2)$. Set

$$R = \mathbf{F}_2[s, t][tX]$$

where X is an indeterminate and set $M = (s, t, tX)$.

Then the following assertions hold:

- (1) M is a maximal ideal of R .
- (2) $|R/M| = 2$.
- (3) (s) is a prime ideal of R .
- (4) $(0) = (s) \cap (t, tX)$.
- (5) (t, tX) is an M -primary ideal of R .
- (6) $\text{Ass}_R(R) = \{(s), M\}$.
- (7) $|R| = \infty$.
- (8) $\overline{\chi}(R) < \infty$.



Proof (1) and (2) are trivial since R/M is isomorphic to \mathbf{F}_2 .

(3) It is clear that $R/(s) \cong \mathbf{F}_2[T][TX]$ and $\mathbf{F}_2[T][TX]$ is an integral domain.

(4) Every element f of R is written as the following form:

$$a_0 + \sum_{n \geq 1} a_n (tX)^n$$

where the sum $\sum_{n \geq 1}$ is a finite sum, a_0 and a_n are in $\mathbf{F}_2[s, t]$ for $n = 1, 2, \dots$. Furthermore, a_0 is written as

$$a_0 = b_0 + b_1 s + \sum_{m \geq 1} c_m t^m$$

and $b_0, b_1, c_m \in \mathbf{F}_2$ for $m = 1, 2, \dots$. Assume that f is in $(s) \cap (t, tX)$. Then $b_0 = b_1 = 0$ because $f \in (t, tX)$, and $a_n = c_m = 0$ for $n = 1, 2, \dots$ and $m = 1, 2, \dots$ because $f \in (s) = \{0, s\}$. Hence $f = 0$. This implies that $(0) = (s) \cap (t, tX)$.

(5) Since $M^2 = (t^2, (tX)^2) \subset (t, tX) \subset M$ and M is maximal, (t, tX) is M -primary by [3, Ch. 4, section 2, no 1, example 3]).

(6) Let $\text{Ann}_R(t)$ be the annihilator ideal of t . Then $(s) = \text{Ann}_R(t)$. Similarly, we have $M = \text{Ann}_R(s)$. Hence $\text{Ass}_R(R) = \{(s), M\}$.

(7) Trivial.

(8) By Proposition 5 below, we see that $\overline{\chi}(R) < \infty$. \square

We call Q a maximal prime divisor of (0) if Q is in $\text{Ass}_R(R)$ and Q is a maximal element in $\text{Ass}_R(R)$ with respect to inclusion. We know that $Z(R) = \cup Q$ where the union is taken over all maximal prime divisors Q 's of (0) in $\text{Ass}_R(R)$. ([4, Theorem 80 and its proof])

Proposition 5 Assume that there is only one maximal element in $\text{Ass}_R(R)$, say M . Then the following assertions hold:

(1) $\overline{\chi}(R)$ is finite if and only if $|R/M| < \infty$.

(2) If $\overline{\chi}(R)$ is finite, then $\overline{\chi}(R) = |R/M|$.

Proof (1) Assume that $\overline{\chi}(R)$ is finite. Let $x_1 + M, x_2 + M, \dots, x_t + M$ be distinct residue classes of R/M . Then $x_i - x_j \notin M = Z(R)$ for $i \neq j$. Therefore, $\{x_1, x_2, \dots, x_t\}$ is a clique of $\overline{G}(R)$. By Lemma 1, $t \leq \overline{\chi}(R)$ and hence $|R/M|$ is finite.

Conversely, assume that $|R/M| < \infty$. Set $|R/M| = t$ and let $x_1 + M, x_2 + M, \dots, x_t + M$ be all residue classes of R/M . Note that $(x + a) - (x + b) = a - b \in M$ for elements $x + a, x + b$ in $x + M$. Set

$$V_1 = x_1 + M, V_2 = x_2 + M, \dots, V_t = x_t + M.$$

Then, we see that

$$V(\overline{G}(R)) = V_1 \coprod V_2 \coprod \dots \coprod V_t.$$

is a disjoint union of $V(\overline{G}(R))$ such that no pair of two distinct elements of V_i is joined by an edge for $i = 1, 2, \dots, t$. Hence by Lemma 2, we get $\overline{\chi}(R) \leq t$. This shows that $\overline{\chi}(R)$ is finite.

(2) From the argument of the proof of the assertion (1), it is easily seen that $\overline{\chi}(R) = |R/M|$. \square

Lemma 6 Let P_1, \dots, P_t be distinct prime ideals of R . Let n be a positive integer and assume that

$$|R/P_1| \geq n, \dots, |R/P_t| \geq n.$$

Then there exist elements z_1, \dots, z_n of R such that

$$z_i - z_j \notin P_1 \cup \dots \cup P_t$$

for $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$.



Proof Let

$$x_u^{(1)} + P_u, x_u^{(2)} + P_u, \dots, x_u^{(n)} + P_u$$

be distinct residue classes of R/P_u for $u = 1, 2, \dots, t$. Set $S = R - P_1 \cup \dots \cup P_t$ and $I = P_1 \cap \dots \cap P_t$. Then R_S is a semi-local ring with maximal ideals $(P_1)_S, \dots, (P_t)_S$. By Chinese remainder theorem we have an isomorphism

$$\phi : R_S/I_S \longrightarrow R_S/(P_1)_S \times \dots \times R_S/(P_t)_S$$

by

$$\phi\left(\frac{x}{s} + I_S\right) = \left(\frac{x}{s} + (P_1)_S, \dots, \frac{x}{s} + (P_t)_S\right).$$

Set

$$\begin{aligned} \alpha_1 &= (x_1^{(1)} + (P_1)_S, x_2^{(1)} + (P_2)_S, \dots, x_t^{(1)} + (P_t)_S), \\ &\dots \\ \alpha_n &= (x_1^{(n)} + (P_1)_S, x_2^{(n)} + (P_2)_S, \dots, x_t^{(n)} + (P_t)_S). \end{aligned}$$

Then there exist elements $\frac{y_1}{s_1}, \dots, \frac{y_n}{s_n}$ of R_S ; ($y_1, \dots, y_n \in R$ and $s_1, \dots, s_n \in S$) such that

$$\phi\left(\frac{y_1}{s_1} + I_S\right) = \alpha_1, \dots, \phi\left(\frac{y_n}{s_n} + I_S\right) = \alpha_n.$$

Set $s = s_1 \cdot \dots \cdot s_n$ and

$$z_1 = s \frac{y_1}{s_1}, \dots, z_n = s \frac{y_n}{s_n}.$$

Then z_1, \dots, z_n are elements of R and

$$\phi(z_1 + I_S) = s\alpha_1, \dots, \phi(z_n + I_S) = s\alpha_n.$$

We shall show that $z_i - z_j \notin P_1 \cup \dots \cup P_t$ for $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$. Suppose the contrary. Then there are indices i, j with $i \neq j$ and u such that $z_i - z_j \in P_u$. Note that $x_u^{(i)} - x_u^{(j)} \notin P_u$, $z_i - sx_u^{(i)} \in (P_u)_S$ and $z_j - sx_u^{(j)} \in (P_u)_S$. Then $z_i - z_j - (sx_u^{(i)} - sx_u^{(j)})$ is in $(P_u)_S$. Since $z_i - z_j \in P_u$, we get $s(x_u^{(i)} - x_u^{(j)}) \in (P_u)_S$. This implies that $x_u^{(i)} - x_u^{(j)} \in P_u$ because $s \notin P_u$. This is a contradiction. \square

Proposition 7 Let R be a Noetherian ring and let P_1, \dots, P_t be all maximal prime divisors of (0) . If $\overline{\chi}(R)$ is finite, then there exists an index u such that $|R/P_u|$ is finite.

Proof Assume that $\overline{\chi}(R)$ is finite and

$$|R/P_1| = \infty, \dots, |R/P_t| = \infty.$$

Set $n = \overline{\chi}(R) + 1$. Then by Lemma 6 there exist elements z_1, \dots, z_n of R such that

$$z_i - z_j \notin P_1 \cup \dots \cup P_t = Z(R)$$

for $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$. Hence $\{z_1, \dots, z_n\}$ is a clique of $\overline{G}(R)$, and by Lemma 1 we have $n \leq \overline{\chi}(R) = n - 1$. This is absurd. Therefore, there exists an index u such that $|R/P_u|$ is finite. \square

Lemma 8 Let P be a prime ideal of R . Let G' be the graph whose vertices are elements of R and in which two distinct vertices x and y are joined by an edge if $x - y \notin P$. If $|R/P|$ is finite, then $\chi(G') = |R/P|$.



Proof Set $n = |R/P|$ and let $x_1 + P, \dots, x_n + P$ be distinct residue classes of R/P . Set $V_1 = x_1 + P, \dots, V_n = x_n + P$. Then

$$V(G') = V_1 \coprod \dots \coprod V_n$$

is a disjoint union of $V(G')$ such that no pair of two distinct elements of V_i is joined by an edge for $i = 1, 2, \dots, n$. By Lemma 2, we get $\chi(G') \leq n$. On the other hand, $\{x_1, \dots, x_n\}$ is a clique of G' , hence $\chi(G') \geq n$. Therefore, $\chi(G') = |R/P|$. \square

Lemma 9 *Let R be a Noetherian ring and let P be a maximal prime divisor of (0) . If $|R/P|$ is finite, then $\overline{\chi}(R)$ is finite and $\overline{\chi}(R) \leq |R/P|$.*

Proof Let G' be the graph as in Lemma 8. Since $P \subset Z(R)$, we see that $\overline{G}(R)$ is a subgraph of G' . Lemma 3 implies that $\overline{\chi}(R) \leq \chi(G')$. By Lemma 8, we have $\chi(G') = |R/P|$. Hence, $\overline{\chi}(R)$ is finite and $\overline{\chi}(R) \leq |R/P|$. \square

Theorem 10 *Let R be a Noetherian ring. Then the following assertions hold:*

- (1) $\overline{\chi}(R)$ is finite if and only if there exists an element P of $\text{Ass}_R(R)$ such that $|R/P|$ is finite.
- (2) Let P_1, \dots, P_t be all maximal prime divisors of (0) . If $\overline{\chi}(R)$ is finite, then

$$\overline{\chi}(R) = \min\{|R/P_i|; i = 1, 2, \dots, t\}.$$

Proof (1) If $\overline{\chi}(R)$ is finite, then by Proposition 7 there exists a maximal prime divisor P of (0) such that $|R/P|$ is finite.

Conversely, if there exists an element P of $\text{Ass}_R(R)$ such that $|R/P|$ is finite. Then there exists a maximal prime divisor Q of (0) such that $P \subset Q$. Since $|R/Q| \leq |R/P|$, we see that $|R/Q|$ is finite. Then, Lemma 9 shows that $\overline{\chi}(R)$ is finite.

- (2) If $\overline{\chi}(R)$ is finite, then $\min\{|R/P_i|; i = 1, 2, \dots, t\}$ is finite by the argument of the proof of the assertion (1). Set $n = \min\{|R/P_i|; i = 1, 2, \dots, t\}$. Then $|R/P_1| \geq n, \dots, |R/P_t| \geq n$. By Lemma 6, there exist elements z_1, \dots, z_n of R such that $z_i - z_j \notin P_1 \cup \dots \cup P_t = Z(R)$ for $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$. Hence $\{z_1, \dots, z_n\}$ is a clique of $\overline{G}(R)$ and $\overline{\chi}(R) \geq n$ by Lemma 1. On the other hand, by Lemma 9, $\overline{\chi}(R) \leq n$. Hence

$$\overline{\chi}(R) = n = \min\{|R/P_i|; i = 1, 2, \dots, t\}.$$

\square

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Atiyah, M.F.; MacDonald, I.G.: Introduction to Commutative Algebra. Addison-Wesley Publishing Com. Inc. (1969)
2. Bondy, J.A.; Murthy, V.S.R.: Graph Theory With Applications. Macmillan Press Ltd. (1976)
3. Bourbaki, N.: Éléments De Mathématique. Algèbre commutative, Hermann (1961)
4. Kaplansky, I.: Commutative Rings, 2nd edn. The University of Chicago Press (1974)
5. Matsumura, H.: Commutative Algebra, 2nd edn. Benjamin/Cummings Publishing Company (1980)
6. Sato, J.; Baba, K.: The chromatic number of the simple graph associated with a commutative ring. *Scientiae Mathematicae Japonicae*, **71**(2), 187–194 (2010)

